



# basic education

---

Department:  
Basic Education  
**REPUBLIC OF SOUTH AFRICA**

**SENIOR CERTIFICATE EXAMINATIONS/  
SENIORSERTIFIKAAT-EKSAMEN  
NATIONAL SENIOR CERTIFICATE EXAMINATIONS/  
NASIONALE SENIORSERTIFIKAAT-EKSAMEN**

**MATHEMATICS P1/WISKUNDE VI**

**MARKING GUIDELINES/NASIENRIGLYNE**

**2022**

**MARKS: 150  
PUNTE: 150**

**These marking guidelines consist of 16 pages.  
*Hierdie nasienriglyne bestaan uit 16 bladsye.***


**NOTE:**

- If a candidate answers a question TWICE, only mark the FIRST attempt.
- Consistent Accuracy applies in all aspects of the marking guidelines.

**LET WEL:**

- Indien 'n kandidaat 'n vraag TWEE keer beantwoord, merk slegs die EERSTE poging.
- Volgehoue akkuraatheid is DEURGAANS op ALLE aspekte van die nasienriglyne van toepassing.

**QUESTION/VRAAG 1**

1.1.1	$x^2 + 2x - 15 = 0$ $(x + 5)(x - 3) = 0$ $x = -5 \text{ or } x = 3$	✓ factors ✓ $x = -5$ ✓ $x = 3$ (3)
1.1.2	$5x^2 - x - 9 = 0$ $x = \frac{-(-1) \pm \sqrt{(-1)^2 - 4(5)(-9)}}{2(5)}$ $x = \frac{1 \pm \sqrt{181}}{10}$ $x = 1,45 \text{ or } x = -1,25$	✓ substitution into the correct formula  ✓ $x = 1,45$ ✓ $x = -1,25$ (3)
1.1.3	$x^2 \leq 3x$ $x^2 - 3x \leq 0$ $x(x - 3) \leq 0$  $0 \leq x \leq 3 \text{ OR } x \in [0;3]$	✓ standard form ✓ factors  ✓✓ answer (4)
1.2.1	$a + \frac{64}{a} = 16$ $a^2 - 16a + 64 = 0$ $(a - 8)^2 = 0$ $a = 8$	✓ standard form ✓ factors  ✓ answer (3)

<p>1.2.2</p>	$2^x + 2^{6-x} = 16$ $2^x + \frac{64}{2^x} = 16$ $2^x = 8 \text{ (from 1.2.1)}$ $2^x = 2^3$ $x = 3$	<p>✓ exp law</p> <p>✓ <math>2^x = 8</math></p> <p>✓ answer (3)</p>
<p>1.3</p>	$\sqrt{\frac{2^{1002}(1+2^4)}{17(2)^{998}}}$ $= \sqrt{\frac{2^4(17)}{17}}$ $= \sqrt{2^4}$ $= 2^2$ $= 4$	<p>✓ common factor</p> <p>✓ second factor</p> <p>✓ simplification</p> <p>✓ answer (4)</p>
<p>1.4</p>	$2x - y = 2 \quad \dots(1)$ $\frac{1}{x} - 3y = 1 \quad \dots(2)$ $y = 2x - 2$ $\frac{1}{x} - 3(2x - 2) = 1$ $\frac{1}{x} - 6x + 6 - 1 = 0$ $1 - 6x^2 + 6x - x = 0$ $-6x^2 + 5x + 1 = 0$ $6x^2 - 5x - 1 = 0$ $(6x + 1)(x - 1) = 0$ $x = -\frac{1}{6} \text{ or } x = 1$ $y = 2\left(-\frac{1}{6}\right) - 2 \text{ or } y = 2(1) - 2$ $y = -\frac{7}{3} \text{ or } y = 0$	<p>✓ <math>y = 2x - 2</math></p> <p>✓ substitution</p> <p>✓ simplification</p> <p>✓ standard form</p> <p>✓ x-values</p> <p>✓ y-values (6)</p>

**OR/OF**

$$x = \frac{2+y}{2} \dots(1)$$

$$\frac{1}{x} - 3y = 1 \dots(2)$$

$$\frac{1}{\frac{2+y}{2}} - 3y = 1$$

$$\frac{2}{2+y} - 3y = 1$$

$$\frac{2 - 6y - 3y^2}{2+y} = 1$$

$$2 - 6y - 3y^2 = 2 + y$$

$$-3y^2 - 7y = 0$$

$$-y(3y + 7) = 0$$

$$y = 0 \quad \text{or} \quad y = -\frac{7}{3}$$

$$x = 1 \quad \text{or} \quad x = -\frac{1}{6}$$

**OR/OF**

$$\checkmark x = \frac{2+y}{2}$$

✓ substitution

✓ simplification

✓ standard form

✓ y-values

✓ x-values

(6)  
[26]

**QUESTION/VRAAG 2**

<p>2.1.1</p>	$a + 6d = 35$ $-1 + 6d = 35$ $6d = 36$ $d = 6$ <p><b>OR/OF</b></p> $\frac{35 - (-1)}{7 - 1} = 6$ <div style="border: 1px solid black; padding: 5px; width: fit-content; margin: 10px auto;"> <p>ANSWER ONLY: FULL MARKS</p> </div>	<p>✓ substitution</p> <p>✓ answer (2)</p> <p><b>OR/OF</b></p> <p>✓ substitution</p> <p>✓ answer (2)</p>
<p>2.1.2</p>	$T_n = a + (n - 1)d$ $473 = -1 + (n - 1)(6)$ $79 = n - 1$ $\therefore n = 80$ <div style="border: 1px solid black; padding: 5px; width: fit-content; margin: 10px auto;"> <p>ANSWER ONLY: FULL MARKS</p> </div>	<p>✓ substitution into the correct formula</p> <p>✓ equating to 473</p> <p>✓ answer (3)</p>
<p>2.1.3</p>	$S_n = \frac{n}{2}[2a + (n - 1)d]$ $S_{40} = \frac{40}{2}[2(-1) + (40 - 1)(6)]$ $\therefore S_{40} = 4640$ <p><b>OR/OF</b></p> $T_{40} = 6(40) - 7$ $= 233$ $S_n = \frac{n}{2}(a + l)$ $= \frac{40}{2}(-1 + 233)$ $= 4640$	<p>✓ substitution</p> <p>✓ answer (2)</p> <p><b>OR/OF</b></p> <p>✓ substitution</p> <p>✓ answer (2)</p>
<p>2.2.1</p>	$  \begin{array}{ccccccc}  75 & & 53 & & 35 & & 21 \\  & \backslash & / & \backslash & / & \backslash & / \\  & -22 & & -18 & & -14 & \\  & & \backslash & / & \backslash & / & \\  & & 4 & & 4 & &   \end{array}  $ <p><math>T_5 = 11</math></p>	<p>✓ answer (A) (1)</p>
<p>2.2.2</p>	$T_n = an^2 + bn + c$ $2a = 4$ $a = 2$ $3a + b = -22$ $6 + b = -22$ $b = -28$ $a + b + c = 75$ $2 - 28 + c = 75$ $c = 101$ $\therefore T_n = 2n^2 - 28n + 101$	<p>✓ <math>T_n = an^2 + bn + c</math></p> <p>✓ <math>a = 2</math></p> <p>✓ <math>b = -28</math></p> <p>✓ <math>c = 101</math> (4)</p>

2.2.3

Minimum value of  $T_n$

$$n = -\frac{b}{2a} = -\frac{(-28)}{2(2)}$$

$$n = 7$$

✓  $n = 7$

Minimum value of  $T_n = 2(7)^2 - 28(7) + 101 = 3$

✓ min value = 3

Each term in the new pattern is  $-\frac{1}{5}$  the value of the terms in the old pattern.

✓  $-\frac{1}{5}$  value of term of old pattern

Maximum value of new pattern =  $-\frac{3}{5}$

✓ max value =  $-\frac{3}{5}$  (4)

**OR/OF**

$$T'_n = 4n - 28$$

$$4n - 28 = 0$$

$$4n = 28$$

$$n = 7$$

**OR/OF**

✓  $n = 7$

Minimum value of  $T_n = 2(7)^2 - 28(7) + 101 = 3$

✓ min value = 3

Each term in the new pattern is  $-\frac{1}{5}$  the value of the terms in the old pattern.

✓  $-\frac{1}{5}$  value of term of old pattern

Maximum value of new pattern =  $-\frac{3}{5}$

✓ max value =  $-\frac{3}{5}$  (4)

**OR/OF**

$$T_n = -\frac{2}{5}n^2 + \frac{28}{5}n - \frac{101}{5}$$

$$n = -\frac{b}{2a} = \frac{-\frac{28}{5}}{2\left(\frac{-2}{5}\right)}$$

$$= 7$$

$$T_7 = -\frac{3}{5}$$

**OR/OF**

✓✓  $T_n \div (-5)$

✓  $n = 7$

✓ max value =  $-\frac{3}{5}$  (4)

**OR/OF**

**OR/OF**

$$T_n = -\frac{2}{5}n^2 + \frac{28}{5}n - \frac{101}{5}$$

$$T'_n = -\frac{4}{5}n + \frac{28}{5}$$

$$\checkmark\checkmark T_n \div (-5)$$

$$-\frac{4}{5}n + \frac{28}{5} = 0$$

$$-4n = -28$$

$$n = 7$$

$$\text{Minimum value of } T_n = 2(7)^2 - 28(7) + 101 = 3$$

Each term in the new pattern is  $-\frac{1}{5}$  the value of the terms in the old pattern.

$$\checkmark n = 7$$

$$\text{Maximum value of new pattern} = -\frac{3}{5}$$

$$\checkmark \text{max value} = -\frac{3}{5}$$

(4)

**[16]**

**QUESTION/VRAAG 3**

<p>3.1.1</p>	$T_n = ar^{n-1}$ $T_{10} = 1024 \left(\frac{1}{4}\right)^{10-1}$ $\therefore T_{10} = \frac{1}{256}$ <div style="border: 1px solid black; padding: 5px; width: fit-content; margin: 10px auto;"> <p style="text-align: center;">ANSWER ONLY: FULL MARKS</p> </div>	<p>✓ substitution into the correct formula</p> <p>✓ answer</p> <p style="text-align: right;">(2)</p>
<p>3.1.2</p>	$\sum_{p=0}^8 256(4^{1-p}) = 1024 + 256 + 64 + \dots$ $S_n = \frac{a[1-r^n]}{1-r}$ $S_9 = \frac{1024 \left[ 1 - \left(\frac{1}{4}\right)^9 \right]}{1 - \frac{1}{4}}$ $S_9 = \frac{87381}{64}$ $= 1365,33$ <p><b>OR/OF</b></p> $\sum_{p=0}^8 256(4^{1-p})$ $= 1024 + 256 + 64 + 16 + 4 + 1 + \frac{1}{4} + \frac{1}{16} + \frac{1}{64}$ $S_9 = \frac{87381}{64}$ $= 1365,33$	<p>✓ 1024</p> <p>✓ <math>n = 9</math></p> <p>✓ substitution into the correct formula</p> <p>✓ answer</p> <p style="text-align: right;">(4)</p> <p><b>OR/OF</b></p> <p>✓ 1024</p> <p>✓ rest of expansion</p> <p>✓ <math>n = 9</math> terms</p> <p>✓ answer</p> <p style="text-align: right;">(4)</p>
<p>3.2</p>	$-t^2 - 6t - 9; \frac{t^3 + 9t^2 + 27t + 27}{2}$ $-(t^2 + 6t + 9); \frac{1}{2}(t+3)(t^2 + 6t + 9)$ $-(t+3)^2; \frac{1}{2}(t+3)^3$ $r = \frac{-(t+3)}{2}$ $-1 < \frac{-t-3}{2} < 1$ $-2 < -t-3 < 2$ $1 < -t < 5$ $-5 < t < -1$	$r = \frac{t^3 + 9t^2 + 27t + 27}{-t^2 - 6t - 9}$ <p>✓ <math>-(t^2 + 6t + 9)</math></p> <p>✓ <math>\frac{1}{2}(t+3)(t^2 + 6t + 9)</math></p> <p>✓ <math>-1 &lt; \frac{-t-3}{2} &lt; 1</math></p> <p>✓ answer</p> <p style="text-align: right;">(5)</p>
		<p><b>[11]</b></p>



**QUESTION 4**

4.1	$10 = a\left(\frac{1}{3}\right)^{-2} + 7$ $3 = 9a$ $\therefore a = \frac{1}{3}$	✓ subs (-2 ; 10) ✓ simplification ✓ answer (3)
4.2	$y = g(0)$ $y = \frac{1}{3} \times \left(\frac{1}{3}\right)^0 + 7$ $y = \frac{22}{3} = 7,33$ $\therefore \left(0 ; \frac{22}{3}\right)$ <div style="border: 1px solid black; padding: 5px; display: inline-block; margin-left: 100px;">ANSWER ONLY: FULL MARKS</div>	✓ substitution of $x = 0$ ✓ answer (2)
4.3.1	Translation by 1 unit to the right and 7 units downwards	✓ 1 unit right ✓ 7 units downwards (2)
4.3.2	$h(x) = \left(\frac{1}{3}\right)^x$ $h^{-1}: x = \left(\frac{1}{3}\right)^y$ $y = \log_{\frac{1}{3}}(x) \quad \text{OR/OF} \quad y = -\log_3(x)$ <div style="border: 1px solid black; padding: 5px; display: inline-block; margin-left: 100px;">ANSWER ONLY: FULL MARKS</div>	✓ swap $x$ and $y$ ✓ answer (2)
		<b>[9]</b>

## QUESTION 5

5.1	$g(x) = \frac{a}{x+2} + q$ Subs (1 ; 0): $0 = \frac{a}{1+2} + q$ $0 = a + 3q$ Subs $\left(0 ; -\frac{1}{2}\right)$ $-\frac{1}{2} = \frac{a}{0+2} + q$ $-1 = a + 2q$ Solving simultaneously: $q = 1$ $a = -3$ $\therefore g(x) = \frac{-3}{x+2} + 1$	$\checkmark g(x) = \frac{a}{x+2} + q$  $\checkmark 0 = a + 3q$  $\checkmark -1 = a + 2q$ $\checkmark \text{ solving simultaneously}$ $\checkmark q = 1$ $\checkmark a = -3$ <p style="text-align: right;">(6)</p>
5.2	$y \in \mathbb{R}; y \neq 1$ <b>OR/OF</b> $(-\infty; 1) \text{ or } (1; \infty)$ <b>OR/OF</b> $y < 1 \text{ or } y > 1$	$\checkmark \text{ answer}$ <p style="text-align: right;">(1)</p>
5.3	$y - 1 = 1(x + 2)$ $y = x + 3$ <div style="border: 1px solid black; padding: 5px; display: inline-block; margin-left: 100px;"> <b>ANSWER ONLY: FULL MARKS</b> </div>	$1 = 1(-2) + c$ $c = 3$ $y = x + 3$ $\checkmark m = 1$ $\checkmark \text{ subs point } (-2 ; 1)$ $\checkmark \text{ answer}$ <p style="text-align: right;">(3)</p>
5.4	$K' (-3 ; 4)$	$\checkmark \text{ x-value}$ $\checkmark \text{ y-value}$ <p style="text-align: right;">(2)</p>
		<b>[12]</b>

**QUESTION 6**

<p>6.1</p>	$f(x) = -x^2 - 6x + 7$ $f'(x) = -2x - 6$ $-2x - 6 = 0$ <p style="text-align: center;"><b>OR/OF</b></p> $x = -\frac{(-6)}{2(-1)}$ $x = -3$ <p>E(-3 ; 16)</p> <div style="border: 1px solid black; padding: 5px; width: fit-content; margin: 10px auto;"> <p style="text-align: center;">ANSWER ONLY: FULL MARKS</p> </div>	<p>✓ method</p> <p>✓ x-value</p> <p>✓ y-value</p> <p style="text-align: right;">(3)</p>
<p>6.2</p>	$k = f(-5)$ $k = -(-5)^2 - 6(-5) + 7$ $\therefore k = 12$	<p>✓ answer (A)</p> <p style="text-align: right;">(1)</p>
<p>6.3</p>	<p>C(0 ; 7)</p> <p>D(-5 ; 12)</p> $m_{CD} = \frac{12 - 7}{-5 - 0}$ $m_{CD} = -1$ <p>Equation of CD:</p> $y = -x + 7$	<p>✓ coordinates of C</p> <p>✓ substitution</p> <p>✓ m</p> <p>✓ answer</p> <p style="text-align: right;">(4)</p>
<p>6.4</p>	$-2x - 6 = -1$ $-2x = 5$ $x = -\frac{5}{2}$ $y = f\left(-\frac{5}{2}\right) = -\left(-\frac{5}{2}\right)^2 - 6\left(-\frac{5}{2}\right) + 7 = \frac{63}{4} = 15,75$ $\therefore P\left(-\frac{5}{2}; \frac{63}{4}\right)$	<p>✓ <math>f'(x) = -2x - 6</math></p> <p>✓ equating to -1</p> <p>✓ x-value</p> <p>✓ y-value (A)</p> <p style="text-align: right;">(4)</p>
<p>6.5</p>	<p>Point by symmetry: (-1 ; 12)</p> $-5 < x < -1$ <p style="text-align: center;"><b>OR/OF</b></p> $-x^2 - 6x + 7 > 12$ $-x^2 - 6x - 5 > 0$ $x^2 + 6x + 5 < 0$ $(x + 1)(x + 5) < 0$ $-5 < x < -1$ <div style="border: 1px solid black; padding: 5px; width: fit-content; margin: 10px auto;"> <p style="text-align: center;">ANSWER ONLY: FULL MARKS</p> </div>	<p>✓ -1</p> <p>✓ answer</p> <p style="text-align: right;">(2)</p> <p>✓ -1</p> <p>✓ answer</p> <p style="text-align: right;">(2)</p>
		<p><b>[14]</b></p>

**QUESTION 7**

<p>7.1</p>	$A = P(1 + i)^n$ $2 = 1 \left( 1 + \frac{0,085}{4} \right)^{4n}$ $4n = \log_{\left( 1 + \frac{0,085}{4} \right)} 2$ $n = 8,24 \text{ years}$	<p> <math>\checkmark 2</math> } In  <math>\checkmark \frac{0,085}{4}</math> } correct                      formula   <math>\checkmark</math> use of logs   <math>\checkmark</math> answer in years                       (4)                 </p>
<p>7.2.1</p>	$A = P(1 - i)^n$ $180\,000 = 500\,000(1 - i)^5$ $\frac{9}{25} = (1 - i)^5$ $\sqrt[5]{\frac{9}{25}} = 1 - i$ $i = 0,1848068\dots$ $r = 18,48\%$	<p> <math>\checkmark</math> subs into correct                      formula   <math>\checkmark</math> simplification   <math>\checkmark i = 0,1848\dots</math>   <math>\checkmark</math> answer                       (4)                 </p>
<p>7.2.2</p>	$A = P(1 + i)^n$ $A = 500\,000(1 + 0,063)^5$ $A = R678\,635,11$	<p> <math>\checkmark</math> subs into correct                      formula  <math>\checkmark</math> answer                       (2)                 </p>
<p>7.2.3</p>	<p>Sinking Fund = 678 635,11 – 180 000 = R 498 635,11</p> $498\,635,11 = \frac{x \left[ \left( 1 + \frac{0,1025}{12} \right)^{58} - 1 \right] \left( 1 + \frac{0,1025}{12} \right)^3}{\frac{0,1025}{12}}$ $x = R6\,510,36$	<p> <math>\checkmark</math> value of sinking fund   <math>\checkmark \frac{0,1025}{12}</math>  <math>\checkmark n = 58</math> (A)  <math>\checkmark \left( 1 + \frac{0,1025}{12} \right)^3</math>  <math>\checkmark</math> answer (A)                       (5)                 </p>
		<p><b>[15]</b></p>

**QUESTION/VRAAG 8**


<p>8.1</p>	$f(x) = -x^2$ $f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$ $f'(x) = \lim_{h \rightarrow 0} \frac{-(x+h)^2 + x^2}{h}$ $f'(x) = \lim_{h \rightarrow 0} \frac{-x^2 - 2xh - h^2 + x^2}{h}$ $= \lim_{h \rightarrow 0} \frac{-2xh - h^2}{h}$ $= \lim_{h \rightarrow 0} \frac{h(-2x - h)}{h}$ $= \lim_{h \rightarrow 0} (-2x - h)$ $\therefore f'(x) = -2x$ <p><b>OR/OF</b></p> $f(x) = -x^2$ $f(x+h) = -(x+h)^2 = -x^2 - 2xh - h^2$ $f(x+h) - f(x) = -x^2 - 2xh - h^2 - (-x^2) = -2xh - h^2$ $f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$ $= \lim_{h \rightarrow 0} \frac{-2xh - h^2}{h}$ $= \lim_{h \rightarrow 0} \frac{h(-2x - h)}{h}$ $= \lim_{h \rightarrow 0} (-2x - h)$ $\therefore f'(x) = -2x$	<p>✓ substitution into formula</p> <p>✓ <math>-(x^2 + 2xh + h^2)</math></p> <p>✓ <math>-2xh - h^2</math></p> <p>✓ <math>-2x - h</math></p> <p>✓ answer (5)</p> <p><b>OR/OF</b></p> <p>✓ <math>-x^2 - 2xh - h^2</math></p> <p>✓ <math>-2xh - h^2</math></p> <p>✓ substitution into the formula</p> <p>✓ <math>-2x - h</math></p> <p>✓ answer (5)</p>
<p>8.2.1</p>	$f(x) = 4x^3 - 5x^2$ $f'(x) = 12x^2 - 10x$	<p>✓ <math>12x^2</math> (A)</p> <p>✓ <math>-10x</math> (A)</p> <p>(2)</p>
<p>8.2.2</p>	$D_x \left[ \frac{-6\sqrt[3]{x} + 2}{x^4} \right]$ $= D_x \left[ \frac{-6(x)^{\frac{1}{3}}}{x^4} + \frac{2}{x^4} \right]$ $= D_x \left[ -6x^{-\frac{11}{3}} + 2x^{-4} \right]$ $= 22x^{-\frac{14}{3}} - 8x^{-5}$	<p>✓ <math>x^{\frac{1}{3}}</math></p> <p>✓ <math>-6x^{-\frac{11}{3}} + 2x^{-4}</math></p> <p>✓ <math>22x^{-\frac{14}{3}}</math></p> <p>✓ <math>-8x^{-5}</math></p> <p>(4)</p>
		<p><b>[11]</b></p>

**QUESTION/VRAAG 9**

<p>9.1</p>	$f(x) = (x+t)^2(x-3)$ $-3 = (0+t)^2(0-3)$ $1 = t^2$ $t = \pm 1$ $\therefore t = 1$ $f(x) = (x+1)^2(x-3)$ $f(x) = (x^2 + 2x + 1)(x-3)$ $f(x) = x^3 - x^2 - 5x - 3$	<p>✓ <math>f(x) = (x+t)^2(x-3)</math>                  ✓ subs (0 ; -3)</p> <p>✓ <math>t</math></p> <p>✓ <math>f(x) = (x+1)^2(x-3)</math>                  ✓ expansion</p> <p style="text-align: right;">(5)</p>
<p>9.2</p>	$f'(x) = 3x^2 - 2x - 5$ $0 = 3x^2 - 2x - 5$ $0 = (x+1)(3x-5)$ $x = -1 \text{ or } x = \frac{5}{3}$ $N\left(\frac{5}{3}; -\frac{256}{27}\right) = (1,67; -9,48)$	<p>✓ <math>f'(x) = 3x^2 - 2x - 5</math>                  ✓ = 0</p> <p>✓ factors                  ✓ x-value (<math>x &gt; 0</math>)</p> <p>✓ y-value (A) (5)</p>
<p>9.3.1</p>	<p><math>x &lt; 3 ; x \neq -1</math></p> <p><b>OR/OF</b>  <math>x &lt; -1 \text{ or } -1 &lt; x &lt; 3</math></p> <p><b>OR/OF</b>  <math>(-\infty; -1) \text{ or } (-1; 3)</math></p>	<p>✓ <math>x &lt; 3</math>                  ✓ <math>x \neq -1</math> (2)</p> <p><b>OR/OF</b>                  ✓ <math>x &lt; -1</math>                  ✓ <math>-1 &lt; x &lt; 3</math> (2)</p> <p><b>OR/OF</b>                  ✓ <math>(-\infty; -1)</math>                  ✓ <math>(-1; 3)</math> (2)</p>
<p>9.3.2</p>	<p><math>x &lt; -1 \text{ or } x &gt; \frac{5}{3}</math> OR/OF <math>x \leq -1 \text{ or } x \geq \frac{5}{3}</math></p> <p><b>OR/OF</b>  <math>(-\infty; -1) \text{ or } \left(\frac{5}{3}; \infty\right)</math> OR/OF <math>(-\infty; -1] \text{ or } \left[\frac{5}{3}; \infty\right)</math></p>	<p>✓ <math>x &lt; -1</math>                  ✓ <math>x &gt; \frac{5}{3}</math> (2)</p> <p><b>OR/OF</b>                  ✓ <math>(-\infty; -1)</math>                  ✓ <math>\left(\frac{5}{3}; \infty\right)</math> (2)</p>
<p>9.3.3</p>	<p><math>f''(x) &gt; 0</math>  <math>6x - 2 &gt; 0</math>  <math>x &gt; \frac{1}{3} \text{ or } \left(\frac{1}{3}; \infty\right)</math></p> <p><b>OR/OF</b>  <math>\frac{\frac{5}{3} + (-1)}{2} = \frac{1}{3}</math>  <math>x &gt; \frac{1}{3} \text{ or } \left(\frac{1}{3}; \infty\right)</math></p> <div style="border: 1px solid black; padding: 5px; width: fit-content; margin: 10px auto;"> <p style="text-align: center;">ANSWER ONLY: FULL MARKS</p> </div>	<p>✓ <math>6x - 2</math>                  ✓ <math>\frac{1}{3}</math>                  ✓ <math>x &gt; \frac{1}{3}</math> (3)</p> <p><b>OR/OF</b>                  ✓ substitution                  ✓ <math>\frac{1}{3}</math>                  ✓ <math>x &gt; \frac{1}{3}</math> (3)</p>

9.4	$\text{Distance} = x^3 - x^2 - 5x - 3 - (3x^2 - 2x - 5)$ $= x^3 - 4x^2 - 3x + 2$ $\frac{d\text{Distance}}{dx} = 3x^2 - 8x - 3$ $0 = 3x^2 - 8x - 3$ $0 = (3x + 1)(x - 3)$ $x = 3 \text{ or } x = -\frac{1}{3}$ <p>Max distance</p> $= \left(-\frac{1}{3}\right)^3 - 4\left(-\frac{1}{3}\right)^2 - 3\left(-\frac{1}{3}\right) + 2$ $= \frac{68}{27} = 2,52$	$\checkmark x^3 - 4x^2 - 3x + 2$ $\checkmark \frac{d\text{Distance}}{dx} = 3x^2 - 8x - 3$ $\checkmark$ factors $\checkmark$ x-values $\checkmark x = -\frac{1}{3}$ $\checkmark$ answer
		(6) <b>[23]</b>

**QUESTION/VRAAG 10**

10.1.1	$7! = 5\ 040$	✓✓ answer (2)
10.1.2	$4! \times 4!$ $= 576$ $P(\text{African flags together}) = \frac{576}{5040} \left( = \frac{4}{35} = 0,11 \right)$	✓ 4! ✓ $4! \times 4!$ ✓ answer (A) (3)
10.2	$P(A \text{ or } B) = P(A) + P(B) - P(A \text{ and } B)$ $0,88 = 0,4 + P(B) - P(A \text{ and } B)$ $0,88 = 0,4 + P(B) - 0,4P(B)$ $0,48 = 0,6P(B)$ $P(B) = 0,8$	✓ subs into rule ✓ $P(A \text{ and } B) = 0,4P(B)$ ✓ answer (3)
10.3	<p style="text-align: center;">                 First Passenger                  Second Passenger             </p>  <p>                 Probability of first passenger choosing meat = <math>\frac{x}{120}</math>                  Probability of second passenger choosing cheese = <math>\frac{120-x}{119}</math>  <math>\frac{x}{120} \times \frac{120-x}{119} = \frac{18}{85}</math>  <math>120x - x^2 = 3\ 024</math>  <math>x^2 - 120x + 3\ 024 = 0</math>  <math>(x - 84)(x - 36) = 0</math>  <math>x = 84 \text{ or } x = 36</math>  <math>\therefore P(\text{1st cheese}) = \frac{36}{120} = \frac{3}{10}</math> </p>	✓ $\frac{x}{120}$ ✓ $\frac{120-x}{119}$ ✓ $\frac{x}{120} \times \frac{120-x}{119} = \frac{18}{85}$ ✓ $x = 84 \text{ or } x = 36$ ✓ $\frac{3}{10}$ (5)
		<b>[13]</b>

**TOTAL/TOTAAL: 150**