



education
MPUMALANGA PROVINCE
REPUBLIC OF SOUTH AFRICA

NATIONAL
SENIOR CERTIFICATE

GRADE 12

PHYSICAL SCIENCES: CHEMISTRY P2

SEPTEMBER 2023

Stanmorephysics

MARKS: 150

TIME: 3 hours

This question paper consists of 15 pages and 4 data sheets.

QUESTION 1: MULTIPLE-CHOICE QUESTIONS

Four options are provided as possible answers to the following questions. Each question has only ONE correct answer. Choose the answer and write only the letter (A–D) next to the question number (1.1 – 1.10) in the ANSWER BOOK, for example, 1.11 E.

- 1.1 Which ONE of the following organic condensed structures is 2,3-dimethylpentane?
- A $(CH_3)_3CCH(CH_3)_2$
 - B $(CH_3)_2CHCH_2CH(CH_3)_2$
 - C $CH_3CH_2C(CH_3)_2CH_2CH_3$
 - D $(CH_3)_2CHCH(CH_3)CH_2CH_3$ (2)
- 1.2 The boiling points of branched alkanes are lower than those of straight chain alkanes containing the same number of carbon atoms because branched alkane chains have ...
- A larger molecular mass.
 - B longer chain lengths.
 - C smaller effective molecular surface areas.
 - D more electrons. (2)
- 1.3 Consider the reaction given below.



Which ONE of the following combinations correctly identifies the type of reaction that takes place and the IUPAC name of product Z?


	TYPE OF REACTION	Z
A	Addition	Propane
B	Addition	Propene
C	Elimination	Propane
D	Elimination	Propene



(2)

- 1.4 Consider the graphs **A** and **B** below. Graph **A** shows the potential energy curve for the same reaction.

Graph **B** shows the distribution of molecular energy. **X** represents the minimum kinetic energy needed for a reaction to take place.




Which of the following will be the correct value for **X** in graph **B**?

- A 103 kJ
- B 93 kJ
- C 35 kJ
- D 10 kJ

(2)

- 1.5 Consider the following hypothetical reaction taking place in a container that has a fixed volume:



The temperature of the system is doubled. Which ONE of the following combinations correctly indicates the change in the NUMBER OF MOLES of $A_3(g)$ and the TOTAL MASS in the container?


	NUMBER OF MOLES OF $A_3(g)$	TOTAL MASS IN CONTAINER
A	Decreases	Increases
B	Increases	Decreases
C	Increases	Remains constant
D	Decreases	Remains constant

(2)

- 1.6 The following reversible reaction reaches equilibrium in a closed container:




The equilibrium was first established after 5 minutes. (The broken line on the graph represents the reverse reaction.)



What possible change could have been made to the reaction conditions at $t = 10$ minutes?

- A The concentration of H_2 increases
 - B The temperature was increased
 - C The temperature was decreased
 - D The external pressure on the reaction mixture was increased. (2)
- 1.7 Chlorine gas, $\text{Cl}_2(\text{g})$, is used to disinfect water in public swimming pools. $\text{Cl}_2(\text{g})$ reacts with water, $\text{H}_2\text{O}(\ell)$ according to the following balanced equation.



The addition of $\text{Cl}_2(\text{g})$ changes the pH of water in the swimming pools.


Which ONE of the following substances must be added to public swimming pools periodically to increase the pH?



- A Na_2CO_3
- B NH_4Cl
- C H_2SO_4
- D KCl (2)

- 1.8 The following equations represent two hypothetical half-reactions.

Which ONE of the following substances from these hypothetical half-reactions will be the strongest oxidising agent?



- A X⁻
- B X₂
- C Y⁺
- D Y

(2)

- 1.9 Which of the following combinations CORRECTLY shows the products formed during the electrolysis of concentrated sodium chloride?

	ANODE	CATHODE
A	Chlorine	Hydrogen
B	Hydrogen	Oxygen
C	Chlorine	Oxygen
D	Hydrogen	Chlorine

(2)

- 1.10 Which ONE of the following equations is spontaneous?

- A Mg²⁺(aq) + Zn(s) → Zn²⁺(aq) + Mg(s)
- B Mg²⁺(aq) + Cu(s) → Cu²⁺(aq) + Mg(s)
- C Cu²⁺(aq) + 2Ag(s) → 2Ag⁺(aq) + Cu(s)
- D 2Ag⁺(aq) + Pb(s) → Pb²⁺(aq) + 2Ag(s)



(2)

[20]



QUESTION 2 (Start on a new page.)

Compound **A** and **B** are hydrocarbons.



- 2.1 Define the underlined term. (1)
- 2.2 Write down the:
 - 2.2.1 IUPAC name of compound **A**. (3)
 - 2.2.2 GENERAL FORMULA of the homologous series to which compound **A** belongs. (1)
 - 2.2.3 The STRUCTURAL FORMULA of compound **B** (3)
 - 2.2.4 NAME of the solution that can be used in the laboratory to test whether compound **A** and **B** are saturated or unsaturated. (1)
- 2.3 Consider the following compound: $\text{CH}_3\text{CH}_2\text{CH}_2\text{CHO}$

Write down the:

 - 2.3.1 Homologous series to which this compound belongs? (1)
 - 2.3.2 NAME of the functional group. (1)
 - 2.3.3 IUPAC name of the CHAIN isomer for this compound. (2)
 - 2.3.4 STRUCTURAL FORMULA of the FUNCTIONAL isomer for this compound. (2)
- 2.4 Consider the compound below:

$(\text{CH}_3)_2\text{C}(\text{OH})\text{CH}_2\text{CH}_2\text{CH}_3$

 - 2.4.1 Write down the IUPAC name of this compound. (2)
 - 2.4.2 Is the compound a PRIMARY, SECONDARY OR TERTIARY alcohol. Give a reason for the answer. (2)



QUESTION 3 (Start on a new page.)

Four compounds (**A** to **D**) are used to investigate factors affecting the melting points. The results obtained are shown in the table below.




	COMPOUNDS	MELTING POINT (°C)
A	Propane	-188
B	Butane	-138
C	Butanal	-96,9
D	Butan-1-ol	-89,8

- 3.1 Define the term *melting point*. (2)
 - 3.2 The melting points of compounds **A** and **B** are compared.
 - 3.2.1 Write down the independent variable for this investigation. (1)
 - 3.2.2 Explain the difference in the melting points of these two compounds. (3)
 - 3.3 Which ONE of the compounds **B**, **C** or **D** has the lowest vapour pressure?
Explain the answer (2)
 - 3.4 The melting points of compounds **C** and **D** are now compared.
 - 3.4.1 Which variable must be kept constant in this experiment? (1)
 - 3.4.2 Refer to the TYPES of intermolecular forces to explain the difference in melting points between compound **C** and **D**. (4)
- [13]



QUESTION 4 (Start on a new page.)

Study the flow diagram of organic reactions below and answer the questions that follow. The letters **X**, **Y** and **Z** represent organic compounds and letters **A** to **E** represent organic reactions.



- 4.1 Write down the type of addition reactions that take place in:
- 4.1.1 Reaction A (1)
 - 4.1.2 Reaction B (1)
 - 4.1.3 Reaction C (1)
- 4.2 For reaction D, write down:
- 4.2.1 The type of reaction that takes place. (1)
 - 4.2.2 TWO reaction conditions necessary for the reaction to occur. (2)
 - 4.2.3 The INORGANIC product that is formed. (1)
- 4.3 Give the FORMULA of the catalyst required for reaction B (1)
- 4.4 Write down a balanced chemical equation using CONDENSED STRUCTURAL FORMULAE for reaction A. Compound X is the main product. (3)
- 4.5 To which homologous series does compound Y belong? (1)
- 4.6 For reaction E, write down the:
- 4.6.1 NAME of the type of reaction (1)
 - 4.6.2 IUPAC name of compound Z. (2)

[15]


QUESTION 5 (Start on a new page.)

The reaction between zinc, $\text{Zn}(\text{s})$, with EXCESS dilute hydrochloric acid, $\text{HCl}(\text{aq})$, is used to investigate factors that affect the rate of a reaction.

The balanced equation for the reaction is:



The apparatus used is illustrated below.



5.1 Define the term *reaction rate*. (2)


5.2 What is the function of the cotton plug? (1)

A summary of the conditions used in EXPERIMENT 1 and 2 is given in the table below.

EXPERIMENT	MASS (Zn)	VOLUME HCl (cm ³)	TEMPERATURE (°C)	STATE OF DIVISION
1	x	150	30	Granules
2	x	150	30	Powder



The change in mass of zinc is calculated and recorded in 3-minute intervals for both experiments. The results obtained are shown in the graph below (NOT drawn to scale).




5.3 Use the information in the graph to answer the following questions:

5.3.1 Which graph **A** or **B** represents EXPERIMENT 2? (1)

5.3.2 Calculate the number of moles of hydrochloric acid that reacted from $t = 3$ minutes to $t = 12$ minutes in the experiment that is represented by graph **A**. (4)

5.4 Calculate the initial mass of zinc used if the average rate of formation of hydrogen gas, in the experiment represented by graph **B**, is $2.5 \times 10^{-4} \text{ mol}\cdot\text{s}^{-1}$. (5)

5.5 A third experiment (EXPERIMENT 3) is carried out by changing only one condition in EXPERIMENT 1. Two energy distribution curves for the reaction in EXPERIMENTS 1 and 3 are shown in the graph below.




5.5.1 What change was made to the reaction conditions in EXPERIMENT 1 to obtain the results of EXPERIMENT 3? (1)

5.5.2 Use the collision theory to explain how the change mentioned in QUESTION 5.5.1 affects the reaction rate. (4)

[18]

QUESTION 6 (Start on a new page.)

A certain amount of sulphur and 0,3 mol oxygen gas are sealed in a container at 340°C. The reaction reaches EQUILIBRIUM according to the following balanced equation:



- 6.1 Define the term *chemical equilibrium*. (2)

- 6.2 How will each of the following changes affect the yield of $SO_2(g)$?

Write down only INCREASES, DECREASES or REMAINS THE SAME.


- 6.2.1 More Sulphur is added to the container. (1)

- 6.2.2 The pressure is increased by decreasing the volume of the container at constant temperature. (1)

- 6.3 It is found that the equilibrium constant (K_c) increases when the temperature is increased. Is the forward reaction EXOTHERMIC or ENDOTHERMIC? Fully explain the answer. (3)

- 6.4 How will the addition of a catalyst influence the equilibrium constant (K_c) of this reaction? Write down only INCREASES, DECREASES or REMAINS THE SAME. (1)

- 6.5 The graph below, not drawn to scale, shows how the amount of $S(s)$ in the container changes with time at 340 °C.



- 6.5.1 Equilibrium is reached at 340 °C. Calculate the equilibrium constant, K_c , at this temperature. The volume of the container is 500 cm³. (9)

- 6.5.2 Determine the K_c value for the reverse reaction. (2)

[19]

QUESTION 7 (Start on a new page.)

- 7.1 The dissociation constant of some substances is given in the table below:



Name of substance	Formula	K _a (298 K)
Hydrogen sulphate ion	HSO ₄ ⁻	1,2 × 10 ⁻²
Ammonium ion	NH ₄ ⁺	5,6 × 10 ⁻¹⁰
Hydrocyanic acid	HCN	4,9 × 10 ⁻¹⁰

- 7.1.1 Write down the NAME or FORMULA of the substance that has the highest tendency to dissociate. Give a reason for the answer. (2)
- 7.1.2 Write down the FORMULAE of the conjugate bases of hydrogen sulphate ion and hydrocyanic acid. (2)
- 7.2 To determine the percentage purity of a sample of oxalic acid (H₂C₂O₄). The three steps below are followed:


Step 1: A standard solution of sodium hydroxide, NaOH(aq), is prepared by adding 50 cm³ of NaOH(aq) of concentration 0,63 mol·dm⁻³ in 950 cm³ of water.

Step 2: An impure sample of oxalic acid solution, H₂C₂O₄(aq), is prepared by dissolving 0,25 g in 75 cm³ of water.

Step 3: The H₂C₂O₄(aq) solution is titrated with the standard NaOH(aq).

During the titration, 40 cm³ of the NaOH(aq) is needed to neutralise 50 cm³ of the impure H₂C₂O₄(aq).


The equation for the reaction is:



- 7.2.1 Define the *endpoint of a titration*. (2)
- 7.2.2 Calculate the concentration of the standard NaOH solution. (3)
- 7.2.3 Calculate the percentage purity of the oxalic acid sample. (6)
- 7.3 Sodium ethanoate solid, CH₃COONa(s), undergoes hydrolysis in solution.
- 7.3.1 Define the term *hydrolysis*. (2)
- 7.3.2 How will the pH of the solution be affected by the hydrolysis of CH₃COONa(s)? Choose from INCREASES, DECREASES or REMAINS THE SAME. Use a balanced equation to explain the answer. (4)
- [21]

QUESTION 8 (Start on a new page.)

The reaction between magnesium and copper(II)nitrate is used to set up an electrochemical cell under standard conditions. A balanced equation of the reaction is given below:




- 8.1 Write down the
- 8.1.1 Type of electrochemical cell (1)
 - 8.1.2 Cell notation of this cell (3)
 - 8.1.3 NAME of the reducing agent (1)
 - 8.1.4 Standard conditions for this electrochemical cell (2)
- 8.2 Calculate the initial emf of this cell. (4)
- 8.3 How will the concentration of Mg²⁺ ions be affected when the cell is functioning? Write down only INCREASES, DECREASES or REMAINS THE SAME. (1)
- 8.4 This cell is connected to a light bulb. In theory the bulb should light up, but in practice it does not. Give TWO possible reasons for this observation. (2)
[14]



QUESTION 9 (Start on a new page.)

The diagram below shows an electrochemical cell used in the refining of copper.



- 9.1 Define the term *electrolyte*. (2)
- 9.2 Write down the formula of the cation in the electrolyte of the above electrochemical cell. (1)
- 9.3 When an electric current passes through the electrolyte the mass of the electrodes changes.
 - 9.3.1 Does the mass of electrode X increase or decrease? (1)
 - 9.3.2 Write down the relevant half reaction to support the answer in QUESTION 9.3.1 (2)
- 9.4 During the process illustrated by the above cell, a total of $2,259 \times 10^{24}$ electrons is transferred. Calculate the mass by which the cathode change. (5)
[11]

TOTAL [150]

DATA FOR PHYSICAL SCIENCES GRADE 12
PAPER 2 (CHEMISTRY)

GEGEWENS VIR FISIESE WETENSKAPPE GRAAD 12
VRAESTEL 2 (CHEMIE)



TABLE 1: PHYSICAL CONSTANTS/TABEL 1: FISIESE KONSTANTES

NAME/NAAM	SYMBOL/SIMBOOL	VALUE/WAARDE
Standard pressure Standaarddruk	p^\ominus	$1,013 \times 10^5 \text{ Pa}$
Molar gas volume at STP Molêre gasvolume by STD	V_m	$22,4 \text{ dm}^3 \cdot \text{mol}^{-1}$
Standard temperature Standaardtemperatuur	T^\ominus	273 K
Avogadro's constant	N_A	$6,023 \times 10^{23} \text{ mol}^{-1}$

TABLE 2: FORMULAE/TABEL 2: FORMULES

$n = \frac{m}{M}$	$n = \frac{N}{N_A}$
$c = \frac{n}{V}$ OR/OF $c = \frac{m}{MV}$	$n = \frac{V}{V_m}$
$\frac{n_a}{n_b} = \frac{c_a V_a}{c_b V_b}$	$\text{pH} = -\log[\text{H}_3\text{O}^+]$
$K_w = [\text{H}_3\text{O}^+][\text{OH}^-] = 1 \times 10^{-14} \text{ at/ by } 298 \text{ K}$	
$E_{\text{cell}}^\ominus = E_{\text{cathode}}^\ominus - E_{\text{anode}}^\ominus / E_{\text{sel}}^\ominus = E_{\text{katode}}^\ominus - E_{\text{anode}}^\ominus$	
OR/OF	
$E_{\text{cell}}^\ominus = E_{\text{reduction}}^\ominus - E_{\text{oxidation}}^\ominus / E_{\text{sel}}^\ominus = E_{\text{reduksie}}^\ominus - E_{\text{oksidasie}}^\ominus$	
OR/OF	
$E_{\text{cell}}^\ominus = E_{\text{oxidising agent}}^\ominus - E_{\text{reducing agent}}^\ominus / E_{\text{sel}}^\ominus = E_{\text{oksideermiddel}}^\ominus - E_{\text{reduseermiddel}}^\ominus$	
$n = \frac{Q}{e}$	$n = \frac{Q}{q_e}$



TABLE 3: THE PERIODIC TABLE OF ELEMENTS

TABLE 4A: STANDARD REDUCTION POTENTIALS
TABEL 4A: STANDAARD-REDUKSIEPOTENSIALE




Increasing strength of oxidising agents/Toenemende sterkte van oksideermiddels

Half-reactions/Halfreaksies		E^θ (V)
F ₂ (g) + 2e ⁻	↑↓	+ 2,87
Co ³⁺ + e ⁻	↑↓	+ 1,81
H ₂ O ₂ + 2H ⁺ + 2e ⁻	↑↓	+ 1,77
MnO ₄ ⁻ + 8H ⁺ + 5e ⁻	↑↓	+ 1,51
Cl ₂ (g) + 2e ⁻	↑↓	+ 1,36
Cr ₂ O ₇ ²⁻ + 14H ⁺ + 4e ⁻	↑↓	+ 1,33
O ₂ (g) + 4H ⁺ + 4e ⁻	↑↓	+ 1,23
MnO ₂ + 4H ⁺ + 2e ⁻	↑↓	+ 1,23
Pt ²⁺ + 2e ⁻	↑↓	+ 1,20
Br ₂ (ℓ) + 2e ⁻	↑↓	+ 1,07
NO ₃ ⁻ + 4H ⁺ + 3e ⁻	↑↓	+ 0,96
Hg ²⁺ + 2e ⁻	↑↓	+ 0,85
Ag ⁺ + e ⁻	↑↓	+ 0,80
NO ₃ ⁻ + 2H ⁺ + e ⁻	↑↓	+ 0,80
Fe ³⁺ + e ⁻	↑↓	+ 0,77
O ₂ (g) + 2H ⁺ + 2e ⁻	↑↓	+ 0,68
I ₂ + 2e ⁻	↑↓	+ 0,54
Cu ⁺ + e ⁻	↑↓	+ 0,52
SO ₂ + 4H ⁺ + 4e ⁻	↑↓	+ 0,45
2H ₂ O + O ₂ + 4e ⁻	↑↓	+ 0,40
Cu ²⁺ + 2e ⁻	↑↓	+ 0,34
SO ₄ ²⁻ + 4H ⁺ + 2e ⁻	↑↓	+ 0,17
Cu ²⁺ + e ⁻	↑↓	+ 0,16
Sn ⁴⁺ + 2e ⁻	↑↓	+ 0,15
S + 2H ⁺ + 2e ⁻	↑↓	+ 0,14
2H ⁺ + 2e ⁻	↑↓	0,00
Fe ³⁺ + 3e ⁻	↑↓	- 0,06
Pb ²⁺ + 2e ⁻	↑↓	- 0,13
Sn ²⁺ + 2e ⁻	↑↓	- 0,14
Ni ²⁺ + 2e ⁻	↑↓	- 0,27
Co ²⁺ + 2e ⁻	↑↓	- 0,28
Cd ²⁺ + 2e ⁻	↑↓	- 0,40
Cr ³⁺ + e ⁻	↑↓	- 0,41
Fe ²⁺ + 2e ⁻	↑↓	- 0,44
Cr ³⁺ + 3e ⁻	↑↓	- 0,74
Zn ²⁺ + 2e ⁻	↑↓	- 0,76
2H ₂ O + 2e ⁻	↑↓	H ₂ (g) + 2OH ⁻ - 0,83
Cr ²⁺ + 2e ⁻	↑↓	Cr - 0,91
Mn ²⁺ + 2e ⁻	↑↓	Mn - 1,18
Al ³⁺ + 3e ⁻	↑↓	Al ^ℓ - 1,66
Mg ²⁺ + 2e ⁻	↑↓	Mg - 2,36
Na ⁺ + e ⁻	↑↓	Na - 2,71
Ca ²⁺ + 2e ⁻	↑↓	Ca - 2,87
Sr ²⁺ + 2e ⁻	↑↓	Sr - 2,89
Ba ²⁺ + 2e ⁻	↑↓	Ba - 2,90
Cs ⁺ + e ⁻	↑↓	Cs - 2,92
K ⁺ + e ⁻	↑↓	K - 2,93
Li ⁺ + e ⁻	↑↓	Li - 3,05



Increasing strength of reducing agents/Toenemende sterkte van reduseermiddels

TABLE 4B: STANDARD REDUCTION POTENTIALS
TABEL 4B: STANDAARD-REDUKSIEPOTENSIALE



Half-reactions/Halreaksies			E^θ (V)
$\text{Li}^+ + \text{e}^-$	\rightleftharpoons	Li	-3,05
$\text{K}^+ + \text{e}^-$	\rightleftharpoons	K	-2,93
$\text{Cs}^+ + \text{e}^-$	\rightleftharpoons	Cs	-2,92
$\text{Ba}^{2+} + 2\text{e}^-$	\rightleftharpoons	Ba	-2,90
$\text{Sr}^{2+} + 2\text{e}^-$	\rightleftharpoons	Sr	-2,89
$\text{Ca}^{2+} + 2\text{e}^-$	\rightleftharpoons	Ca	-2,87
$\text{Na}^+ + \text{e}^-$	\rightleftharpoons	Na	-2,71
$\text{Mg}^{2+} + 2\text{e}^-$	\rightleftharpoons	Mg	-2,36
$\text{Al}^{3+} + 3\text{e}^-$	\rightleftharpoons	Al	-1,66
$\text{Mn}^{2+} + 2\text{e}^-$	\rightleftharpoons	Mn	-1,18
$\text{Cr}^{2+} + 2\text{e}^-$	\rightleftharpoons	Cr	-0,91
$2\text{H}_2\text{O} + 2\text{e}^-$	\rightleftharpoons	$\text{H}_2(\text{g}) + 2\text{OH}^-$	-0,83
$\text{Zn}^{2+} + 2\text{e}^-$	\rightleftharpoons	Zn	-0,76
$\text{Cr}^{3+} + 3\text{e}^-$	\rightleftharpoons	Cr	-0,74
$\text{Fe}^{2+} + 2\text{e}^-$	\rightleftharpoons	Fe	-0,44
$\text{Cr}^{3+} + \text{e}^-$	\rightleftharpoons	Cr^{2+}	-0,41
$\text{Cd}^{2+} + 2\text{e}^-$	\rightleftharpoons	Cd	-0,40
$\text{Co}^{2+} + 2\text{e}^-$	\rightleftharpoons	Co	-0,28
$\text{Ni}^{2+} + 2\text{e}^-$	\rightleftharpoons	Ni	-0,27
$\text{Sn}^{2+} + 2\text{e}^-$	\rightleftharpoons	Sn	-0,14
$\text{Pb}^{2+} + 2\text{e}^-$	\rightleftharpoons	Pb	-0,13
$\text{Fe}^{3+} + 3\text{e}^-$	\rightleftharpoons	Fe	-0,06
$2\text{H}^+ + 2\text{e}^-$	\rightleftharpoons	$\text{H}_2(\text{g})$	0,00
$\text{S} + 2\text{H}^+ + 2\text{e}^-$	\rightleftharpoons	$\text{H}_2\text{S}(\text{g})$	+0,14
$\text{Sn}^{4+} + 2\text{e}^-$	\rightleftharpoons	Sn^{2+}	+0,15
$\text{Cu}^{2+} + \text{e}^-$	\rightleftharpoons	Cu^+	+0,16
$\text{SO}_4^{2-} + 4\text{H}^+ + 2\text{e}^-$	\rightleftharpoons	$\text{SO}_2(\text{g}) + 2\text{H}_2\text{O}$	+0,17
$\text{Cu}^{2+} + 2\text{e}^-$	\rightleftharpoons	Cu	+0,34
$2\text{H}_2\text{O} + \text{O}_2 + 4\text{e}^-$	\rightleftharpoons	4OH^-	+0,40
$\text{SO}_2 + 4\text{H}^+ + 4\text{e}^-$	\rightleftharpoons	$\text{S} + 2\text{H}_2\text{O}$	+0,45
$\text{Cu}^+ + \text{e}^-$	\rightleftharpoons	Cu	+0,52
$\text{I}_2 + 2\text{e}^-$	\rightleftharpoons	2I^-	+0,54
$\text{O}_2(\text{g}) + 2\text{H}^+ + 2\text{e}^-$	\rightleftharpoons	H_2O_2	+0,68
$\text{Fe}^{3+} + \text{e}^-$	\rightleftharpoons	Fe^{2+}	+0,77
$\text{NO}_3^- + 2\text{H}^+ + \text{e}^-$	\rightleftharpoons	$\text{NO}_2(\text{g}) + \text{H}_2\text{O}$	+0,80
$\text{Ag}^+ + \text{e}^-$	\rightleftharpoons	Ag	+0,80
$\text{Hg}^{2+} + 2\text{e}^-$	\rightleftharpoons	$\text{Hg}(\ell)$	+0,85
$\text{NO}_3^- + 4\text{H}^+ + 3\text{e}^-$	\rightleftharpoons	$\text{NO}(\text{g}) + 2\text{H}_2\text{O}$	+0,96
$\text{Br}_2(\ell) + 2\text{e}^-$	\rightleftharpoons	2Br^-	+1,07
$\text{Pt}^{2+} + 2\text{e}^-$	\rightleftharpoons	Pt	+1,20
$\text{MnO}_2 + 4\text{H}^+ + 2\text{e}^-$	\rightleftharpoons	$\text{Mn}^{2+} + 2\text{H}_2\text{O}$	+1,23
$\text{O}_2(\text{g}) + 4\text{H}^+ + 4\text{e}^-$	\rightleftharpoons	$2\text{H}_2\text{O}$	+1,23
$\text{Cr}_2\text{O}_7^{2-} + 14\text{H}^+ + 4\text{e}^-$	\rightleftharpoons	$2\text{Cr}^{3+} + 7\text{H}_2\text{O}$	+1,33
$\text{Cl}_2(\text{g}) + 2\text{e}^-$	\rightleftharpoons	2Cl^-	+1,36
$\text{MnO}_4^- + 8\text{H}^+ + 5\text{e}^-$	\rightleftharpoons	$\text{Mn}^{2+} + 4\text{H}_2\text{O}$	+1,51
$\text{H}_2\text{O}_2 + 2\text{H}^+ + 2\text{e}^-$	\rightleftharpoons	$2\text{H}_2\text{O}$	+1,77
$\text{Co}^{3+} + \text{e}^-$	\rightleftharpoons	Co^{2+}	+1,81
$\text{F}_2(\text{g}) + 2\text{e}^-$	\rightleftharpoons	2F^-	+2,87



NATIONAL
SENIOR CERTIFICATE

GRADE 12

PHYSICAL SCIENCES: CHEMISTRY P2
SEPTEMBER 2023
MARKING GUIDELINES

MARKS/PUNTE: 150



This memorandum consists of 12 pages.

Hierdie memorandum bestaan uit 12 bladsye.

QUESTION 1 / VRAAG 1

1.1 D ✓✓



1.2 C ✓✓

1.3 D ✓✓

1.4 B ✓✓

1.5 C ✓✓

1.6 D ✓✓

1.7 A ✓✓

1.8 B ✓✓

1.9 A ✓✓

1.10 D ✓✓

[20]

QUESTION 2 / VRAAG 2

2.1 Organic compounds that consist of hydrogen and carbon only. ✓
Organiese verbindings wat slegs uit waterstof en koolstof bestaan.

(1)

2.2.1 2,4-dimethylhex-1-ene/2,4-dimetielheks-1-een

Marking criteria:

- Correct stem i.e. hexene. ✓
- Dimethyl✓
- IUPAC name completely correct including numbering, sequence, hyphens and commas. ✓

Nasienkriteria:

- Korrekte stam d.i. hekseen. ✓
- Dimetiel✓
- IUPAC-naam heeltemal korrek insluitende volgorde, koppeltekens en kommas. ✓


(3)

2.2.2 C_nH_{2n} ✓

(1)



2.2.3



Marking criteria / Nasienriglyne

- Seven C atoms in longest chain (only single bonds between C-atoms) / Sewe C-atome in langste ketting (slegs enkel bindings tussen C-atome) ✓
- Two methyl and one ethyl substituents / Twee metiel en een etiel-substituente. ✓
- Whole structure correct / Hele struktuur korrek ✓

(3)

2.2.4 Bromine (water)/Broom(water)/Br₂ ✓

(1)

2.3.1 Aldehyde/Aldehyied ✓

(1)


2.3.2 Formyl groep/Formielgroep ✓

(1)

2.3.3 2-methyl✓ butanal✓ /2-metielbutanaal

(2)

2.3.4



Marking criteria / Nasienriglyne

- Only functional group correct / Slegs funksionele groep korrek ✓
- Whole structure correct / Hele struktuur korrek

(2)

2.4.1 2-methyl✓ -pentan-2-ol✓ /2-metielpentan-2-ol

(2)

2.4.2 Tertiary (alcohol)/Tertiäre (alkohol) ✓

The C-atom bonded to the hydroxyl/-OH is bonded to three other C-atoms. ✓
Die C-atoom wat aan die hidroksiel/-OH gebind is, is aan drie ander C-atome gebind.

(2)

[19]

QUESTION 3 / VRAAG 3

3.1

Marking criteria/Nasienriglyne

If any one of the underlined key phrases in the **correct context** is omitted, deduct 1 mark/Indien enige van die onderstreepte frase in die **korrekte konteks** uitgelaat is, trek 1 punt af.

The temperature at which the solid and liquid phases of a substance are at equilibrium. ✓✓

Die temperatuur waarby die vaste- en vloeistoffases van 'n stof in ewewig is.

(2)



3.2.1 Chain length / number of C-atoms / Molecular mass ✓
Kettinglengte / getal C-atome/ molekulêre massa

(1)

3.2.2 • **Structure:**

Compound B has a larger surface area/longer chain length/more C atoms/larger molecular mass than compound A✓

• **Intermolecular forces:**

Intermolecular forces in compound B are stronger than those in compound A.✓

• **Energy**

More energy needed to overcome intermolecular forces in compound B than in compound A. ✓

• **Struktuur:**

Verbinding B het 'n groter oppervlakte/langer kettinglengte/meer C-atome/groter molekulêre massa as verbinding A.

• **Intermolekulêre kragte:**

Intermolekulêre kragte in verbinding B is sterker as in verbinding A.

• **Energie:**

Meer energie word benodig om intermolekulêre kragte in verbinding B te oorkom as in verbinding A.

(3)

3.3 D/Butan-1-ol ✓

D has the highest boiling point/ *D het die hoogste kookpunt* ✓

(2)

3.4.1 Molecular mass/number of Carbon and Hydrogen atoms/Molekulêre massa/aantal Koolstof en Waterstof atome ✓

(1)

3.4.2 • In C/ butanal/aldehydes: dipole-dipole forces ✓(in addition to London forces/dispersion forces/induced dipole forces).

• In D/butan-1-ol: Hydrogen bonding. ✓ (in addition to London forces/dispersion forces/induced dipole forces).

• Intermolecular forces in C/butanal are weaker ✓than in D/butan-1-ol OR dipole-dipole forces are weaker than hydrogen bonds OR intermolecular forces in D/butan-1-ol are stronger than in C/butanal.

• More energy needed to overcome/break intermolecular forces in D. ✓

• In C/butanal/aldehiede: dipool-dipoolkragte (tesame met Londonkragte/ dispersiekragte/geïnduseerde dipoolkragte).

• In D/butan-1-ol: Waterstofbinding. (tesame met Londonkragte/ dispersiekragte/geïnduseerde dipoolkragte).

• Intermolekulêre kragte in C/butanaal/aldehied is swakker as in D/butan-1-ol/alkohol

• Intermolekulêre kragte in D/butan-1-ol sterker as in C/butanaal OF dipool-dipoolkragte is swakker as waterstofbindings.

• Meer energie benodig om intermolekulêre kragte te oorkom/breek in D.

(4)

[13]




QUESTION 4/ VRAAG 4

- 4.1.1 Hydrohalogenation/Hydrobromination✓
Hidrohalogenasie/Hidrohalogenering/Hidrobrominasie (1)
- 4.1.2 Hydrogenation/Hidrogenering/Hidrogenasie✓ (1)
- 4.1.3 Hydration/Hidrasie✓ (1)
- 4.2.1 Substitution/Hydrolysis✓
Substitusie/Hidrolise (1)
- 4.2.2 Dilute strong base/NaOH/KOH/LiOH✓ and mildly heated✓
Verdunde sterk basis/ NaOH/KOH/LiOH en matig verhit (2)
- 4.2.3 NaBr/Sodium bromide/Natriumbromied✓ (1)
- 4.3 Pt/Pd/Ni✓ (1)

4.4 Marking criteria/Nasienglyne

- Correct condensed structural formula for but-1-ene ✓
Korrekte gekondenseerde struktuurformule van but-1-een
- Compound X: Bromine atom on correct carbon✓ and whole condensed structural formula correct ✓
Verbinding X: Broom atoom op korrekte C-atoom en die hele gekondenseerde struktuurformule korrek.



NOTE/LET WEL

- Penalise only once for the use of structural formulae or molecular formulae
- *Penaliseer slegs een keer vir die gebruik van struktuurformules of molekulêre formule*

(3)

- 4.5 Alcohol/Alkohol ✓ (1)
- 4.6.1 Esterification/Esterifikasie ✓ (1)
- 4.6.2 Butyl ✓ methanoate✓/Butyletanoaat (2)

[15]



QUESTION 5/ VRAAG 5

5.1

Marking criteria/Nasieneriglyne

If any one of the underlined key phrases in the **correct context** is omitted, deduct 1 mark./Indien enige van die onderstreepte frase in die **korrekte konteks** uitgelaat is, trek 1 punt af.

ANY ONE

- The change in concentration ✓ of reactants or products per unit time✓
- Change in amount/number of moles/volume/mass of products or reactants per (unit) time.
- Amount/number of moles/volume/mass of products formed/reactants used per (unit) time.
- Rate of change in concentration/amount of moles/number of moles/volume/mass. ✓✓ (2 or 0)

ENIGE EEN

- Die verandering in konsentrasie ✓ van reaktante of produkte per eenheid tyd.✓
- Verandering in hoeveelheid/getal mol/volume/massa van produkte of reaktanse per (eenheid) tyd.
- Hoeveelheid/getal mol/volume/massa van produkte gevorm/reaktanse gebruik per (eenheid) tyd.
- Tempo van verandering in konsentrasie/ hoeveelheid mol/getal mol/ volume/massa. ✓ (2 of 0)

(2)

5.2

To make sure that only the H₂(g)/gases escapes./Om seker te maak dat slegs H₂(g)/gasse kan ontsnap. ✓

(1)

5.3.1

Graph/Grafiek B ✓

(1)



5.3.2

Marking criteria/Nasienvriglyne

- Calculate change in $m(\text{Zn})$ or $n(\text{Zn})$ ✓
Bereken verandering in $m(\text{Zn})$ of $n(\text{Zn})$
- Substitute/Vervang $65 \text{ g} \cdot \text{mol}^{-1}$ in $n = \frac{m}{M}$ ✓
- Use mol ratio/Gebruik molverhouding: $n(\text{Zn}) = 2n(\text{HCl})$ ✓
- Final answer/Finale antwoord (0,2 mol) ✓

OPTION 1/OPSIE 1	OPTION 2/OPSIE 2
$\Delta Zn = 7 - 0,5 \checkmark$ $= 6,5 \text{ g}$ $n(\text{Zn}_{\text{used/gebruik}}) = \frac{m}{M}$ $= \frac{6,5}{65} \checkmark$ $= 0,1 \text{ mol Zn}$ $n(\text{HCl}) = 2n(\text{Zn}) \checkmark$ $= 2(0,1) \checkmark$ $= 0,2 \text{ mol} \checkmark$	$n(\text{Zn}_{3s}) = \frac{m}{M} = \frac{7}{65} \checkmark = 0,108 \text{ mol Zn}$ $n(\text{Zn}_{12s}) = \frac{0,5}{65}$ $= 0,0077 \text{ mol Zn}$ $\Delta n(\text{Zn}) = 0,108 - 0,0077 \checkmark$ $= 0,1 \text{ mol Zn}$ $n(\text{HCl}) = 2n(\text{Zn}) \checkmark$ $= 2(0,1)$ $= 0,2 \text{ mol} \checkmark$

(4)

5.4

Marking criteria/Nasienvriglyne

- Substitute/Vervang $2,5 \times 10^{-4}$ in ave rate/gem. $\text{Tempo} = \frac{\Delta n}{\Delta t}$ ✓
- Substitute/Vervang 12×60 (720 s) in ave rate/gem. $\text{Tempo} = \frac{\Delta n}{\Delta t}$ ✓
- Use mol ratio/Gebruik molverhouding: $n(\text{Zn}) = 2n(\text{H}_2)$ ✓
- Substitute/Vervang $65 \text{ g} \cdot \text{mol}^{-1}$ in $n = \frac{m}{M}$ ✓
- Final answer/Finale antwoord: 10,08 g ✓

$$\text{ave rate/gem. tempo} = -\frac{\Delta n}{\Delta t}$$

$$\therefore 2,5 \times 10^{-4} \checkmark = -\frac{\Delta n}{720 - (0)} \checkmark$$

$$\Delta n = -0,18 \text{ mol}$$

$$0 - n_i = -0,18$$

$$n(\text{Zn}) = n(\text{H}_2) = 0,18 \text{ mol} \checkmark$$

$$n(\text{Zn}) = 0,18 \times 65 \checkmark$$

$$= 11,7 \text{ g} \checkmark$$



(5)

5.5.1 Temperature was decreased/ Temperatuur was verlaag ✓

(1)

- 5.5.2 • Decrease in temperature decreases the average kinetic energy/molecules move slower. ✓
Afname in temperatuur verlaag die gemiddelde kinetiese energie/molekule beweeg stadiger.
- Less molecules have enough/sufficient kinetic energy/Less molecules have $E_k \geq E_a$. ✓
Minder molekule het genoeg/voldoende kinetiese energie/Minder molekule het $E_k \geq E_a$.
- Less effective collisions per unit time/second. /Frequency of effective collisions decreases. ✓
Meer effektiewe botsings per eenheidtyd/sekonde./Frekwensie van effektiewe botsings neem toe.
- Reaction rate decreases /Reaksietempo neem af✓

(4)
[18]

QUESTION 6 / VRAAG 6

- 6.1 When the rate of the forward reaction equals the rate of the reverse reaction. ✓✓
Die tempo van die voorwaartse reaksie is gelyk aan die tempo van die terugwaartse reaksie. (2)
- 6.2.1 Reamaine the same / Bly dieselfde ✓ (1)
- 6.2.2 Reamaine the same / Bly dieselfde ✓ (1)
- 6.3 Endothermic / Endotermies ✓
 • A increase in temperature favour the endothermic reaction ✓
'n Verhooging in temperatuur bevoordeel die endotermiese reaksie
 • The forward reaction was favoured ✓
Die voorwaartse reaksie is bevoordeel.
 • The forward reaction is endothermic /*Die voorwaartse reaksie is endotermies.* (3)
- 6.4 Reamaine the same / Bly dieselfde ✓ (1)

6.5.1 Marking criteria / Nasienkriteria:

- Subsitute 32 in formula $n = \frac{m}{M}$ ✓
Vervang 32 in formule $n = \frac{m}{M}$
- Substitute 0,334 in $nS(s)_{\text{initial}}$ and 0,253 in $nS(s)_{\text{final}}$ in table✓
Vervang 0,334 in $nS(s)_{\text{aanvanklik}}$ en 0,253 in $nS(s)_{\text{finaal}}$ in tabel
- Use mol ratio / Gebruik mol verhouding : 1:1:1 ✓
- $n_{\text{Equilibrium}}/ n_{\text{Ewewig}} = n_{\text{Reactants}} / n_{\text{Reactante}_{\text{initial/begin}}} - n_{\text{Reactants}} / n_{\text{Reactante}_{\text{reacted/reageer}}}$ ✓
- $n_{\text{Equilibrium}}/ n_{\text{Ewewig}} = n_{\text{SO}_2}_{\text{initial/begin}} + n_{\text{SO}_2}_{\text{reacted/reageer}}$ ✓
- Divide by the volume (0,5)/ Deel deur die volume (0,5) ✓
- Correct K_c expression / Korrekte K_c uitdrukking ✓
- Correct substitute into K_c expression / Korrekte invervanging in K_c -uitdrukking ✓
- Final answer / Finale antwoord (0,37) ✓



$$n_i = \frac{m}{M}$$

$$= \frac{10,7}{32} \checkmark(a)$$

$$= 0,334 \text{ mol}$$

$$n_f = \frac{m}{M}$$

$$= \frac{8,1}{32}$$

$$= 0,253 \text{ mol}$$



	S(s)	O ₂ (g)	SO ₂ (g)
Initial mol Aanvanklike mol	0,334	0,3	0
Change in mol Verandering in mol $\checkmark(b)$	-0,081	-0,081	+0,081
Equilibrium mol Ewewigs mol	0,253	0,219 $\checkmark(d)$	0,081 $\checkmark(e)$
Equilibrium concentration Ewewigskonstante		0,438	0,162

Ratio $\checkmark(c)$

Divided by/deel
deur 0,5 $\checkmark(f)$

$$K_c = \frac{[\text{SO}_2]}{[\text{O}_2]} \checkmark(g)$$

$$= \frac{0,162}{0,438} \checkmark(h)$$

$$= 0,36986 \checkmark(i)$$

$$= 0,37$$

(9)

6.5.2 $K_c = \frac{1}{0,37}$
 $= 2,704 \checkmark\checkmark$

(2)
[19]



QUESTION 7 / VRAAG 7

- 7.1.1 Hydrogen sulphate ion/Waterstofsultaatioon/ HSO_4^- ✓, Ka value is the highest /
Ka waarde is die hoogste✓ (2)
- 7.1.2 SO_4^{2-} ✓ and/en CN^- ✓ (2)
- 7.2.1 The point where the indicator changes colour. ✓✓
Die punt waar die indicator kleur verander. (2)
- 7.2.2 $c_i V_i = c_f V_f$
 $(0,63)(0,05)✓ = c_f(1)✓$
 $c_f = 0,0315 \text{ mol} \cdot \text{dm}^{-3}✓$ (3)

POSITIVE MARKING FROM QUESTION 7.2.2

Marking criteria / Nasienkriteria:

- Use of formula ($n = c \times v$) and ($m = n \times M$)✓
- Substitute answer in 7.2.2 concentration $n_{\text{NaOH}} = c \times v$ ✓
Vervang antwoord in 7.2.2 as konsentrasie in $n_{\text{NaOH}} = c \times V$
- Ratio / Verhouding $n(\text{NaOH}) : 2 : 1$ ✓
- Substitute / Vervang $90 \text{ g} \cdot \text{mol}^{-1}$ in $m(\text{C}_2\text{H}_2\text{O}_4) = n \times M$ ✓
- Divide $m(\text{C}_2\text{H}_2\text{O}_4)$ by given mass / Deel $m(\text{C}_2\text{H}_2\text{O}_4)$ deur gegewe massa✓
- Answer / Antwoord (22,68%)✓

$$\begin{aligned} n(\text{NaOH}) &= cV \\ &= (0,0315)(0,04)✓ \quad (\text{b}) \\ &= 1,26 \times 10^{-3} \text{ mol} \end{aligned} \quad \checkmark(\text{a})$$


$$\begin{aligned} n(\text{C}_2\text{H}_2\text{O}_4) &= \frac{1}{2}n(\text{NaOH}) \\ &= \frac{1}{2}(1,26 \times 10^{-3}) \\ &= 6,3 \times 10^{-4} \text{ mol} \quad \checkmark \quad (\text{c}) \end{aligned}$$

$$\begin{aligned} m(\text{C}_2\text{H}_2\text{O}_4) &= nM \\ &= (6,3 \times 10^{-4})(90)✓ \quad (\text{d}) \\ &= 0,0567 \text{ g} \end{aligned}$$

$$\begin{aligned} \% \text{ purity} / \text{suiwerheid} &= \frac{0,0567}{0,25} \quad \checkmark \quad (\text{e}) \times 100 \\ &= 22,68\% \quad \checkmark \quad (\text{f}) \end{aligned} \quad (6)$$

- 7.3.1 Reaction of a salt with water / Reaksie van 'n sout met water. ✓✓ (2)

- 7.3.2 Increases / Neem toe ✓




OH^- ions formed causes the solution to become more alkaline therefore pH increases/ OH^- ione wat gevorm word veroorsaak dat oplossing meer alkalis word en dus die pH laat toeneem. ✓

(4)

[21]



QUESTION 8 / VRAAG 8

- 8.1.1 Galvanic cell / Galvaniese sel ✓ (1)
- 8.1.2 Mg / Mg²⁺ ✓ // ✓ Cu²⁺ / Cu ✓ (3)
- 8.1.3 Magnesium ✓ (1)
- 
- 8.1.4 Concentration / Konsentrasie: 1 mol·dm⁻³ ✓
Temperature / Temperatuur: 25°C ✓ (2)
- 8.2 $E^\theta_{\text{cell}} = E^\theta_{\text{cathode}} - E^\theta_{\text{anode}}$ ✓
 $= 0,34 \checkmark - (- 2,36) \checkmark$
 $= 2,7V \checkmark$ (4)
- 
- 8.3 Increases / Neem toe ✓ (1)
- 8.4 • The light bulb is a resistor. / Gloeilamp is 'n weerstand✓
• The potential difference is enough but the current is too small./ Die potensiaalverskil is genoeg maar die stroom is te klein. ✓ (2)
- [14]

QUESTION 9 / VRAAG 9

- 9.1 Is a substance of which the aqueous solution contains ions / substance that dissolves in water to give a solution that conducts electricity. ✓✓
'n Stof waarvan die oplossing ione het / 'n Stof wat in water oplos om 'n oplossing te vorm wat elektriese stroom geleei. (2)
- 9.2 Cu²⁺ ✓ (1)
- 9.3.1 Decreases / Afneem (1)
- 9.3.2 Cu → Cu²⁺ + 2e⁻ ✓✓ (2)

9.4 **Marking criteria / Nasienkriteria:**

- Use Formula / gebruik formule $n = \frac{N}{NA}$ ✓
- Answer / antwoord $n = 3,7525$ mol ✓
- Use mol ratio / Gebruik mol verhouding - 2e⁻: Cu 2:1 ✓
- Substitute 63,5 in formula / Vervang 63,5 in formule $n = \frac{m}{M}$ ✓
- Final answer / Finale antwoord (119,13 g) ✓



$$n = \frac{N}{N_A} \checkmark$$
$$= \frac{2,259 \times 10^{24}}{6,02 \times 10^{23}}$$
$$= 3,7525 \text{ mol } \checkmark$$

$$n(\text{Cu}) = \frac{3,7525}{2} \checkmark$$
$$= 1,876 \text{ mol}$$

$$m(\text{Cu}) = nM$$
$$= (1,876)(63,5) \checkmark$$
$$= 119,13 \text{ g } \checkmark$$

(5)
[11]

GRAND TOTAL/GROOTTOTAAL [150]





ERRATA: PHYSICAL SCIENCES 2023

GRADE 12 PREPARATORY EXAM PAPER 2

QUESTION

2.3.3 2-methyl propanal ✓ / 2-metielpropanaal ✓ (2)

ONLY AFRIKAANS QUESTION PAPER ERRATA:

4.6.2 Butiel✓ metanoaat✓ (2)

$$\begin{aligned} 9.4 \quad n &= \frac{N}{N_A} \checkmark \\ &= \frac{2259 \times 10^{24}}{6,02 \times 10^{23}} \\ &= 3752,49 \text{ mol } \checkmark \end{aligned}$$

$$\begin{aligned} n(\text{Cu}) &= \frac{3752,49}{2} \checkmark \\ &= 1876,246 \text{ mol} \end{aligned}$$

$$\begin{aligned} m(\text{Cu}) &= nM \\ &= (1876,246)(63,5) \checkmark \\ &= 119141,61 \text{ g } \checkmark \end{aligned} \quad (5)$$

